

ECALLANTIDE (KALBITOR®) PRESCRIBER ORDER FORM

Fax completed form, insurance information, and clinical documentation to: **(888) 582-7154**



Patient Name:

Date of Birth:

Address:

Phone:

Height:

inches cm

Weight:

lbs. kg

Clinical Information

Primary Diagnosis Description: Defects in the complement system (hereditary angioedema)

ICD-10 Code: D84.1

Ecallantide (Kalbitor®) Prescription

Ecallantide (Kalbitor®) 30 mg (3 x 10 mg/mL vials) refill as directed x 1 year

Inject a total dose of 30 mg SQ as 3 x 10 mg (1 mL) SQ injections PRN acute HAE attack.

If no response in 45 to 60 minutes after initial dose, an additional total dose of 30 mg SQ may be administered.

If no response in 45 to 60 minutes after the second dose, call the prescriber.

Dispense _____ doses.

Keep _____ doses on-hand at all times.

Ancillary Orders

Anaphylaxis Kit

➔ Required per Option Care Health policy – please complete Anaphylaxis Physician Order (FR-PC-036) provided.

Pre-Medication Orders

Other: _____

Skilled nurse to administer doses SQ as ordered. Refill above ancillary orders as directed x 1 year.

I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber Signature: _____ Date: _____

Prescriber Information

Prescriber Name:

Phone:

Fax:

Address:

NPI:

City, State:

Zip:

Office Contact:

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.

ANAPHYLAXIS PRESCRIBER ORDER FORM

Fax completed form, insurance information, and clinical documentation to:



Patient Name:

Date of Birth:

Address:

Weight:

lbs kg

Anaphylaxis Kit Components

Age Group	Medication	Dose	Instructions
Adults & Pediatrics > 30 kg	Diphenhydramine 1 mL (50 mg/mL) vial #1	25 mg	Slow IV push (if line patent) or IM; may repeat x 1 in 15 min PRN if no improvement
	Epinephrine 1 mL ampule/vial (1 mg/mL) #2	0.3 mg	SQ x 1 dose & repeat x 1 in 5 to 15 min PRN
	or		or
	*Epinephrine 0.3 mg auto-injector 2-pack kit #1	IM x 1 dose & may repeat x 1 in 5 to 15 min PRN	
	Normal saline 500 mL bag #1	500 mL	KVO rate PRN anaphylaxis or over 2 to 4 hr PRN headache rated > 5 on pain scale
Pediatrics 15 to 30 kg	Diphenhydramine 1 mL (50 mg/mL) vial #1	1.25 mg/kg	Slow IV push (if line patent) or IM; may repeat x 1 in 15 min PRN if no improvement
	Epinephrine 1 mL ampule/vial (1 mg/mL) #2	0.15 mg	SQ x 1 dose & repeat x 1 in 5 to 15 min PRN
	or		or
	*Epinephrine 0.15 mg auto-injector 2-pack kit #1	IM x 1 dose & may repeat x 1 in 5 to 15 min PRN	
	Normal saline 250 mL bag #1	250 mL	KVO rate PRN anaphylaxis or over 2 to 4 hr PRN headache rated > 5 on pain scale
Pediatrics < 15 kg	Diphenhydramine 1 mL (50 mg/mL) vial #1	1.25 mg/kg	Slow IV push (if line patent) or IM; may repeat x 1 in 15 min PRN if no improvement
	Epinephrine 1 mL ampule/vial (1 mg/mL) #2	0.01 mg/kg	SQ or IM x 1 dose & repeat x 1 in 5 to 15 min PRN
	Normal saline 250 mL bag #1	250 mL	KVO rate PRN anaphylaxis

Dispense supplies necessary to administer aforementioned medications, including syringes and needles.

*For subcutaneous immune globulin patients, only epinephrine auto-injector 2-pack kits will be dispensed.

Management of Anaphylaxis

General Anaphylaxis

1. **STOP** infusion.
2. Administer emergency meds as ordered.
3. Administer epinephrine as above and repeat dose if necessary.
4. Administer injectable diphenhydramine as above or orally per treatment guidelines.
5. If IV line is in place, infuse normal saline.
6. Initiate CPR (if needed).
7. Call EMS (activate the emergency medical system).
8. Monitor vital signs – elevate legs if hypotensive.
9. Notify prescriber and Option Care Health Director of Nursing and pharmacist.

Subcutaneous Immune Globulin Anaphylaxis – For SEVERE reactions such as wheezing, difficulty in breathing, or swelling of eye lids, lips, or throat.

1. **STOP** the infusion of the medication immediately and remove the needles from the skin.
2. Call 911.
3. Administer epinephrine for one dose as above, repeat dose if necessary.
4. Notify prescriber and Option Care Health Director of Nursing or pharmacist.

Intravenous Immune Globulin Anaphylaxis – If nurse is present in the home, for SEVERE reactions including angioedema, wheezing, difficulty in breathing, or swelling of eye lids, lips, or throat.

1. **STOP** the infusion of the medication immediately. Completely remove the source of the infusate while maintaining venous access.
2. Contact or have caregiver call 911.
3. Administer epinephrine as above and may repeat dose if necessary.
4. Administer *injectable* diphenhydramine (Benadryl®) as above.
5. Monitor and document patient's vital signs, including mental status. If hypotensive, place the patient in supine position with lower extremities elevated or in Trendelenburg position. If breathing difficulty, tilt the patient's head or thrust jaw to relieve airway obstruction.
6. Maintain IV line with normal saline (sodium chloride 0.9%) as above to keep line open until the arrival of a paramedic or ambulance.
7. Contact the prescriber and Option Care Health Director of Nursing or pharmacist.
8. If cardiopulmonary arrest occurs, begin CPR.
9. Monitor and document vital signs every 2 minutes until stable, then every 15 minutes as needed.
10. Remain with patient until paramedics arrive.

When appropriate, nurse shall instruct patient/caregiver about the signs/symptoms of allergic, anaphylactic, and adverse reactions along with the proper use of kit medications. This physician order shall be recognized for the patient's period of treatment and/or up to one year.

I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber Signature:

Date:

Prescriber Information

Prescriber Name:

NPI:

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that do not require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.