Uplizna	[®] (INEBILIZUMAB-CDON) PRES	CRIBER O RDER	Form				
Patient Name:			Date of Birth:				
Address:							
Phone:		ŀ	leight:	🗌 inche	s 🗆 cm	Weight:	🗆 Ibs 🗆 kg
			Information				0
Primary Dia	agnosis Description: Neuromyelitis Opti	ca Spectrum Disor	order (NMOSD) ICD-10 Code: G36.0				
Is this the first dose?			Honatitis	B Status:		te:	
		lue:	· · · · · · · · · · · · · · · · · · ·		Positive Negative		
	PPD (negative) – date:		Active TB				
TB Status:	Last chest x-ray – date:	kon	□ Unknown □ QuantiFERON®TB Gold (negative) - Date				
Past positive TB infection, course taken: QuantiFERON®TB Gold (negative) - Date Uplizna Prescription							
 Option Care Health to initiate services beginning with dose number as indicated below: Dose 1: Infuse 300mg. Administer diluted infusion over approx. 90 minutes at an increasing rate. Use a 0.2 or 0.22 micron in-line filter. Start by infusing 42ml/hr for the first 30 min, followed by a rate of 125ml/hr for the next 30 min. Increase to 333ml/hr until complete. Dose 2: (2 weeks after dose 1) Infuse 300mg over 90 minutes at above increasing rate. Subsequent doses: (every 6 months starting 6 months from the first infusion). Infuse 300mg over 90 min at the increasing rate. Withdraw 10ml from each of three 100mg/10ml vials (total 30ml) and add contents to 250ml of 0.9% sodium chloride ONLY. Prior to every infusion, assess for active infection and delay infusion as appropriate. 							
Ancillary Orders Anaphylaxis Kit							
If this is a 1 st infusion, would you like Option Care Health to provide an anaphylaxis kit with the 1 st dose? Yes No Dosage: Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SUBQ or IM x 1; repeat x 1 in 5 to 15 min PRN. Diphenhydramine 25 mg (> 30 kg) or 1.25 mg/kg (≤ 30 kg) IV or IM; repeat x 1 in 15 min PRN no improvement. O.9% Sodium Chloride 500 mL (>30 kg) or 250 mL (≤ 30 kg) IV at KVO rate PRN anaphylaxis. Pre-Medication Orders Methylprednisolone sodium succinate mg IV given 30 min prior to infusion. Diphenhydramine mg PO 30-60 min prior to infusion. Patient may decline. Acetaminophen mg PO 30-60 min before infusion. Patient may decline. Other:							
		-		l I will he su	inervisina	the nationt's tree	atment
I certify that the use of the indicated treatment is medically necessary, and I will be supervising the patient's treatment.							
Prescriber S	ignature:		Date:				
		Prescribe	er Information				
Prescriber Name: Address:			Phone:			Fax:	
City, State: Zip:		1	Office Contact:				
Fax completed form, insurance information, and clinical do							
CONFIDENTIAL H require authoriza obtained. Unauth or entity to whom employee or age	EALTH INFORMATION: Healthcare information is personal i tion. You are obligated to maintain it in a safe, secure, and c orized re-disclosure or failure to maintain confidentiality co n it is addressed and may contain information that is privileg nt responsible for delivering it to the intended recipient, you please notify us immediately. Brand names are the propert	information related to a per- confidential manner. Re-disc uld subject you to penalties ged and confidential, the disc are hereby notified that an	son's healthcare. It is bein losure of this information described in federal and s closure of which is govern y dissemination, distribut	is prohibited unl state laws. IMPC ed by applicable	ess permitted RTANT WARM law. If the rea	by law or appropriate cus NING: This message is intender of this message is not	tomer/patient authorization is inded for the use of the persor the intended recipient, or the